

FIFTEENTH ANNUAL CONFERENCE

YUCOMAT 2013

Hunguest Hotel Sun Resort Herceg Novi, Montenegro,
September 2-6, 2013
<http://www.mrs-serbia.org.rs>

Programme and The Book of Abstracts

Organised by:
Materials Research Society of Serbia

under the auspices of
**Federation of European Material Societies
and
Materials Research Society**

Title: THE FIFTEENTH ANNUAL CONFERENCE
YUCOMAT 2013
Programme and The Book of Abstracts

Publisher: Materials Research Society of Serbia
Knez Mihailova 35/IV, 11000 Belgrade, Serbia
Phone: +381 11 2185-437; Fax: + 381 11 2185-263
<http://www.mrs-serbia.org.rs>

Editors: Prof. Dr. Dragan P. Uskoković and Prof. Dr. Velimir Radmilović

Technical editor: Aleksandra Stojičić

Cover page: Aleksandra Stojičić and Milica Ševkušić

Copyright © 2013 Materials Research Society of Serbia

Acknowledgments:



Printed in: Biro Konto
Sutorina bb, Igalo – Herceg Novi, Montenegro
Phones: +382-31-670123, 670025, E-mail: bkonto@t-com.me
Circulation: 220 copies. The end of printing: August 2013

P.S.A.12.

PREPARATION AND CHARACTERIZATION Pt CATALYST ON Ru DOPED TIN OXIDE SUPPORT FOR OXYGEN REDUCTION

Lj.M. Gajić-Krstajić¹, N.R. Elezović², B.M. Babić³, V. Radmilović⁴, N.V. Krstajić⁴
¹*Institute of Technical Sciences of SASA, Belgrade, Serbia,* ²*Institute for Multidisciplinary Research, University of Belgrade, Belgrade, Serbia,* ³*Vinča Institute of Nuclear Sciences, Belgrade, Serbia,* ⁴*Faculty of Technology and Metallurgy, University of Belgrade, Belgrade, Serbia*

Platinum nanocatalyst on Ru doped tin oxide support (Ru-SnO₂) was synthesized and characterized as a catalyst for oxygen reduction reaction in 0.5 mol dm⁻³ HClO₄ solution, at 25°C. Ru doped tin oxide support was synthesized by sol-gel procedure. Synthesized support was characterized by BET (Brunauer, Emmett, Teller), X-ray diffraction, HRTEM (high resolution transmission electron microscopy) and cyclic voltammetry techniques. Specific surface area of the support determined from nitrogen adsorption/desorption isothermal curves was 141 m² g⁻¹. XRD spectra contained mainly SnO₂ belonging peaks. The corresponding crystallite size for Ru-SnO₂, determined by Scherrer's equation was 4nm.

Platinum nanocatalyst at Ru-SnO₂ support was synthesized by borohydride reduction method and characterized by X-ray and TEM techniques. Homogenous Pt nanoparticles distribution over the support, without pronounced particle agglomeration was confirmed. The average Pt particle size was 5.3 nm. Electrochemically active surface area of the catalyst was determined from adsorption/desorption charge of hydrogen atoms, after double layer charge correction, taking into account the reference value of 210 μC cm⁻² for full monolayer coverage with adsorbed hydrogen species.

The oxygen reduction reaction at Pt/Ru-SnO₂ catalyst was studied by cyclic voltammetry and linear sweep voltammetry at rotating disc electrode. Two different Tafel slope were observed: one close to 60 mV dec⁻¹ in low current density region, and other close to 120 mV dec⁻¹ at high current densities region, as it was already reported in literature for oxygen reduction at pure polycrystalline Pt, as well as at Pt nanoparticles in acid solutions. The specific activities, expressed in terms of kinetic current densities per electrochemically active surface area, as well as per mass of Pt loaded, at the constant potential of practical interest, were compared to carbon supported (Vulcan XC-72) catalyst. Pt/Ru-SnO₂ catalyst exhibited similar catalytic activity for oxygen reduction reaction like carbon supported one. The durability of the catalysts was evaluated by repetitive cycling up to 1.4 V vs RHE. Better stability of Pt/Ru-SnO₂ catalyst compared to Pt on carbon support was confirmed by determination of the loss of platinum electrochemically active surface area after potential cycling tests.