Supplementary data for the article: P.M. Batinić, V.B. Đorđević, S.I. Stevanović, B.D. Balanč, S.B. Marković, N.D. Luković, D.Ž. Mijin, B.M. Bugarski, Formulation and characterization of novel liposomes containing histidine for encapsulation of a poorly soluble vitamin, Journal of Drug Delivery Science and Technology, 2020, 59, 101920, doi: https://doi.org/10.1016/j.jddst.2020.101920



This work is licensed under the Creative Commons - Attribution 4.0 International (CC BY 4.0) license

Formulation and characterization of novel liposomes containing histidine for encapsulation of a poorly soluble vitamin

Petar M. Batinić¹, Verica B. Đorđević², Sanja I. Stevanović³, Bojana D. Balanč¹, Smilja B. Marković⁴, Nevena D. Luković², Dušan Ž. Mijin^{2,*}, Branko M. Bugarski²

¹Innovation Center of Faculty of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, 11000 Belgrade, Serbia

²Faculty of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, 11000 Belgrade, Serbia

³Institute of Chemistry, Technology and Metallurgy, Department of Electrochemistry, University of Belgrade, Njegoševa 12, 11000 Belgrade, Serbia

⁴Institute of Technical Science of SASA, Knez Mihailova 35/IV, 11000 Belgrade, Serbia




Figure S1. The spectrum of folic acid

^{*}Corresponding author: D. Ž. Mijin (<u>kavur@tmf.bg.ac.rs</u>), (tel: +381113303671, fax: +381113370387)




Figure S2. The spectrum of histidine




Figure S3. The spectrum of phospholipon 90 G




Figure S4. The spectrum of cholesterol




Figure S5. The spectrum of SPAN 20