



ACADEMIE SERBE DES SCIENCES ET DES ARTS

BULLETIN

TOME CLIII

CLASSE DES SCIENCES
MATHÉMATIQUES ET NATURELLES

SCIENCES MATHÉMATIQUES

N° 45

BULLETIN T. CLIII - SCIENCES MATHÉMATIQUES N° 45

BULLETIN

TOME CLIII

CLASSE DES SCIENCES
MATHÉMATIQUES ET NATURELLES

SCIENCES MATHÉMATIQUES

N° 45



BULLETIN

TOME CLIII

CLASSE DES SCIENCES
MATHEMATIQUES ET NATURELLES

SCIENCES MATHEMATIQUES

N° 45

Rédacteur

GRADIMIR V. MILOVANOVIĆ

Membre de l'Académie

BEOGRAD 2020

Publié et imprimé par
Académie serbe des sciences et des arts
Beograd, Knez Mihailova 35

En raison de la pandémie de COVID-19, ce numéro du Bulletin a été publié en 2021

Tirage 300 exemplaires

© Académie serbe des sciences et des arts, 2020

TABLE DES MATIÈRES

1. M. Kostić: <i>On hypercyclicity and supercyclicity of strongly continuous semigroups induced by semiflows. Disjoint hypercyclic semigroups</i>	1
2. E. Malkowsky, V. Rakočević, V. Veličković: <i>Bounded linear and compact operators between the Hahn space and spaces of strongly summable and bounded sequences</i>	25
3. I. Gutman: <i>Sombor index – one year later</i>	43
4. G. V. Milovanović, A. Mir, A. Hussain, A. Ahmad: <i>A note on sharpening of the Erdős-Lax inequality concerning polynomials</i>	57
5. M. M. Mateljević: <i>The Ahlfors-Schwarz lemma, curvature, distance and distortion</i>	67

A NOTE ON SHARPENING OF THE ERDŐS-LAX INEQUALITY CONCERNING POLYNOMIALS

GRADIMIR V. MILOVANOVIĆ, ABDULLAH MIR, ADIL HUSSAIN, ABRAR AHMAD

(Presented at the 5th Meeting, held on June 16, 2021)

A b s t r a c t. In this paper, we establish some upper bound estimates for the modulus of the derivative of a polynomial on the unit disk while taking into account the placement of the zeros and the extremal coefficients of the polynomial. The obtained results sharpen as well as generalize the Erdős-Lax inequality and its extension by Aziz and Dawood and other related inequalities.

AMS Mathematics Subject Classification (2020): 30A10, 30C10, 30C15.

Key Words: Bernstein-type inequalities, Extremal coefficients, Erdős-Lax inequality.

1. Introduction

For an arbitrary entire function f , set $\|f\| = \max_{|z|=1} |f(z)|$, the uniform-norm of f on the unit disk $|z| = 1$. Let $P(z) = \sum_{\nu=0}^n a_{\nu} z^{\nu}$ be an algebraic polynomial of degree n in the complex plane and $P'(z)$ its derivative. The Bernstein-type inequalities relating the uniform-norm of the derivative and the underlying polynomial on the unit circle in the plane play a key role in the literature for proving the inverse theorems in approximation theory, and of course have their own intrinsic interest. These inequalities for constrained polynomials have been the subject of many research papers which is witnessed by many recent articles (for example, see [6], [8, 9], [12]–[15]). A classical result due to Bernstein is that: for two polynomials

$P(z)$ and $Q(z)$ with degree of $P(z)$ not exceeding that of $Q(z)$ and $Q(z) \neq 0$ for $|z| > 1$, the inequality $|P(z)| \leq |Q(z)|$ on the unit circle $|z| = 1$ implies the inequality of their derivatives $|P'(z)| \leq |Q'(z)|$ on $|z| = 1$. In particular, this classical result allows one to establish the famous Bernstein inequality [4] for the uniform-norm on the unit circle. Namely, if $P(z)$ of degree n , it is true that

$$\|P'(z)\| \leq n\|P(z)\|. \quad (1.1)$$

Equality holds in (1.1) if and only if $P(z)$ has all its zeros at the origin. It might easily be observed that the restriction on the zeros of $P(z)$ imply an improvement in (1.1). It turns out that to have any hope of an improved upper bound, one must have some control over the location of the zeros of polynomial $P(z)$. It was conjectured by Erdős and later proved by Lax [10] that if $P(z)$ is a polynomial of degree n and $P(z) \neq 0$ in $|z| < 1$, then

$$\|P'(z)\| \leq \frac{n}{2}\|P(z)\|. \quad (1.2)$$

Inequality (1.2) was further sharpened by Aziz and Dawood [2] in the form of

$$\|P'(z)\| \leq \frac{n}{2} \left\{ \|P(z)\| - \min_{|z|=1} |P(z)| \right\}. \quad (1.3)$$

Equality in (1.2) and (1.3) holds for any polynomial which has all its zeros on $|z| = 1$.

Various extension of (1.2) and (1.3) are known in the literature on various regions of the complex plane (for example, see Govil [5], Malik [11], Milovanović et al.[13] and Rahman and Schmeisser [16]). The purpose of this paper is to generalize and strengthen the inequalities (1.2), (1.3) and related results, while taking into account the placement of the zeros and the extremal coefficients of the polynomial.

2. Main results

We begin by presenting the following extension and refinement of (1.2).

Theorem 2.1. *If $P(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$ is a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$, then for $|z| = 1$,*

$$|P'(z)| \leq \frac{\|P\|}{2} \left[n - \left\{ n \left(\frac{k-1}{k+1} \right) + \frac{2}{k+1} \left(\frac{|a_0| - k^n |a_n|}{|a_0| + k^n |a_n|} \right) \right\} \frac{|P(z)|^2}{\|P\|^2} \right]. \quad (2.1)$$

Equality in (2.1) holds for $P(z) = (z+k)^n$, evaluated at $z = 1$.

Taking $k = 1$ in (2.1), we get the following refinement of (1.2).

Corollary 2.1. *If $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$ is a polynomial of degree n and $P(z) \neq 0$ in $|z| < 1$, then for $|z| = 1$,*

$$|P'(z)| \leq \frac{\|P\|}{2} \left\{ n - \left(\frac{|a_0| - |a_n|}{|a_0| + |a_n|} \right) \frac{|P(z)|^2}{\|P\|^2} \right\}. \quad (2.2)$$

Equality in (2.2) holds for $P(z) = z^n + 1$.

Remark 2.1. In fact, excepting the case when all the zeros of $P(z)$ lie on unit disk $|z| = 1$, the bound obtained in (2.2) is always sharp than the bound obtained in (1.2).

The following result which was recently proved by Ahanger and Shah [1] immediately follows from Theorem 2.1 as a special case.

Corollary 2.2. *If $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$ is a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$, then for $|z| = 1$,*

$$|P'(z)| \leq \frac{n\|P\|}{2} \left\{ 1 - \frac{k-1}{k+1} \cdot \frac{|P(z)|^2}{\|P\|^2} \right\}. \quad (2.3)$$

Equality in (2.3) holds for $P(z) = (z+k)^n$, evaluated at $z = 1$.

Next, we prove the following extension and sharpening of (1.3), which also provides a generalization of Theorem 2.1.

Theorem 2.2. *If $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$ is a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$, then for $0 \leq t \leq 1$ and $|z| = 1$,*

$$|P'(z)| \leq \frac{\|P\| - tm}{2} \left\{ n - \left[\frac{(k-1)n}{k+1} + \frac{2C_n(k,t)}{k+1} \right] \left(\frac{|P(z)| - tm}{\|P\| - tm} \right)^2 \right\}, \quad (2.4)$$

where

$$m = \min_{|z|=k} |P(z)| \quad \text{and} \quad C_n(k,t) = \frac{|a_0| - tm - k^n|a_n|}{|a_0| - tm + k^n|a_n|}.$$

Equality in (2.4) holds for $P(z) = (z+k)^n$, evaluated at $z = 1$.

For $k = 1$, Theorem 2.2 reduces to the following result which is a refinement of (1.3).

Corollary 2.3. *If $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$ is a polynomial of degree n and $P(z) \neq 0$ in $|z| < 1$, then for $0 \leq t \leq 1$ and $|z| = 1$,*

$$|P'(z)| \leq \frac{\|P\| - tm'}{2} \left\{ n - \left(\frac{|a_0| - tm' - |a_n|}{|a_0| - tm' + |a_n|} \right) \left(\frac{|P(z)| - tm'}{\|P\| - tm'} \right)^2 \right\}, \quad (2.5)$$

where

$$m' = \min_{|z|=1} |P(z)|.$$

Equality in (2.5) holds for $P(z) = z^n + 1$.

Remark 2.2. For $t = 0$, (2.4) reduces to (2.1) and (2.5) reduces to (2.2).

The following result which is a refinement of (2.3) immediately follows from Theorem 2.2 as a special case.

Corollary 2.4. *If $P(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$ is a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$, then for $0 \leq t \leq 1$ and $|z| = 1$,*

$$|P'(z)| \leq \frac{n(\|P\| - tm)}{2} \left\{ 1 - \frac{k-1}{k+1} \left(\frac{|P(z)| - tm}{\|P\| - tm} \right)^2 \right\}, \quad (2.6)$$

where

$$m = \min_{|z|=k} |P(z)|.$$

Equality in (2.6) holds for $P(z) = (z+k)^n$, evaluated at $z = 1$.

3. Auxiliary results

For the proofs of the theorems, we shall make use of the following lemmas.

Lemma 3.1. *If x_ν , $\nu = 1, 2, \dots, n$, is a sequence of real numbers such that for each $n \in \mathbb{N}$, $x_\nu \geq 1$, then*

$$(\forall n \in \mathbb{N}) \quad \sum_{\nu=1}^n \frac{x_\nu - 1}{x_\nu + 1} \geq \frac{\prod_{\nu=1}^n x_\nu - 1}{\prod_{\nu=1}^n x_\nu + 1}.$$

PROOF. The above lemma follows by using induction on n . We omit the details.

Lemma 3.2. *If $P(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$ is a polynomial of degree n having no zeros in $|z| < k$, $k \geq 1$, then for each point z on $|z| = 1$ for which $P(z) \neq 0$, we have*

$$\operatorname{Re} \left(\frac{zP'(z)}{P(z)} \right) \leq \frac{1}{1+k} \left\{ n - \frac{|a_0| - k^n|a_n|}{|a_0| + k^n|a_n|} \right\}. \quad (3.1)$$

PROOF. Recall that $P(z)$ has all its zeros in $|z| \geq k, k \geq 1$. If z_1, z_2, \dots, z_n , are the zeros of $P(z) = \sum_{\nu=0}^n a_\nu z^\nu$ of degree n , then $|z_\nu| \geq k, k \geq 1$, and we can write $P(z) = a_n \prod_{\nu=1}^n (z - z_\nu)$. This gives

$$\frac{zP'(z)}{P(z)} = \sum_{\nu=1}^n \frac{z}{z - z_\nu}.$$

Now for the points $e^{i\theta}, 0 \leq \theta \leq 2\pi$, with $P(e^{i\theta}) \neq 0$, we have

$$\begin{aligned} \operatorname{Re} \left(\frac{e^{i\theta} P'(e^{i\theta})}{P(e^{i\theta})} \right) &= \sum_{\nu=1}^n \operatorname{Re} \left(\frac{e^{i\theta}}{e^{i\theta} - z_\nu} \right) \\ &\leq \sum_{\nu=1}^n \frac{1}{1 + |z_\nu|} \\ &= \frac{n}{1+k} - \frac{1}{1+k} \sum_{\nu=1}^n \frac{|z_\nu| - k}{|z_\nu| + 1} \\ &\leq \frac{n}{1+k} - \frac{1}{1+k} \sum_{\nu=1}^n \frac{|z_\nu| - k}{|z_\nu| + k} \quad (\text{as } k \geq 1) \\ &= \frac{n}{1+k} - \frac{1}{1+k} \sum_{\nu=1}^n \frac{\frac{|z_\nu|}{k} - 1}{\frac{|z_\nu|}{k} + 1}. \end{aligned}$$

Since $|z_\nu|/k \geq 1, \nu = 1, 2, \dots, n$, we get on using Lemma 3.1 for the points $e^{i\theta}, 0 \leq \theta \leq 2\pi$, with $P(e^{i\theta}) \neq 0$,

$$\begin{aligned} \operatorname{Re} \left(\frac{e^{i\theta} P'(e^{i\theta})}{P(e^{i\theta})} \right) &\leq \frac{n}{1+k} - \frac{1}{1+k} \left(\frac{\prod_{\nu=1}^n \frac{|z_\nu|}{k} - 1}{\prod_{\nu=1}^n \frac{|z_\nu|}{k} + 1} \right) \\ &= \frac{n}{1+k} - \frac{1}{1+k} \left(\frac{\frac{|a_0|}{k^n |a_n|} - 1}{\frac{|a_0|}{k^n |a_n|} + 1} \right), \end{aligned}$$

which is equivalent to (3.1). This completes the proof of Lemma 2.

Lemma 3.3. *If $P(z)$ is a polynomial of degree n having all its zeros in $|z| \leq k, k \leq 1$, then for $|z| = 1$ and $R > 1$,*

$$|P(Rz) - P(z)| \geq \left(\frac{R^n - 1}{k^n} \right) \min_{|z|=k} |P(z)|. \tag{3.2}$$

The above lemma is due to Aziz and Rather [3].

If we divide both sides of (3.2) by $R - 1$ and let $R \rightarrow 1$, we immediately get the following result.

Lemma 3.4. *If $P(z)$ is a polynomial of degree n having all its zeros in $|z| \leq k$, $k \leq 1$, then for $|z| = 1$, we have*

$$|zP'(z)| \geq \frac{n}{k^n} \min_{|z|=k} |P(z)|.$$

Lemma 3.5. *If $P(z)$ is a polynomial of degree n , then for $|z| = 1$, we have*

$$|P'(z)| + |Q'(z)| \leq n \|P(z)\|,$$

where here and throughout $Q(z) = z^n \overline{P(1/\bar{z})}$.

The above lemma is due to Govil and Rahman [7].

4. Proofs of main results

PROOF OF THEOREM 2.1. Recall that $P(z)$ has no zeros in $|z| < k$, $k \geq 1$. First suppose that $k > 1$, therefore by Lemma 3.2, we have for $|z| = 1$,

$$2 \operatorname{Re} \left(\frac{zP'(z)}{P(z)} \right) \leq \frac{2}{1+k} \left\{ n - \frac{|a_0| - k^n |a_n|}{|a_0| + k^n |a_n|} \right\}. \quad (4.1)$$

If $Q(z) = z^n \overline{P(1/\bar{z})}$, then it easily follows that

$$|Q'(z)| = |nP(z) - zP'(z)| \quad \text{for } |z| = 1.$$

This implies for $|z| = 1$,

$$\begin{aligned} \left| \frac{Q'(z)}{P(z)} \right|^2 &= \left| n - \frac{zP'(z)}{P(z)} \right|^2 \\ &= n^2 + \left| \frac{zP'(z)}{P(z)} \right|^2 - 2n \operatorname{Re} \left(\frac{zP'(z)}{P(z)} \right), \end{aligned}$$

which by (4.1) yields for $|z| = 1$,

$$\left| \frac{zQ'(z)}{P(z)} \right|^2 \geq \left| \frac{zP'(z)}{P(z)} \right|^2 + n^2 \left(\frac{k-1}{k+1} \right) + \frac{2n}{1+k} \left(\frac{|a_0| - k^n |a_n|}{|a_0| + k^n |a_n|} \right).$$

This gives for $|z| = 1$,

$$\left[|P'(z)|^2 + n \left\{ n \frac{k-1}{k+1} + \frac{2}{k+1} \cdot \frac{|a_0| - k^n |a_n|}{|a_0| + k^n |a_n|} \right\} |P(z)|^2 \right]^{1/2} \leq |Q'(z)|. \quad (4.2)$$

Inequality (4.2) gives, by using Lemma 3.5, that for $|z| = 1$,

$$\begin{aligned} |P'(z)|^2 + n \left\{ n \left(\frac{k-1}{k+1} \right) + \frac{2}{k+1} \left(\frac{|a_0| - k^n |a_n|}{|a_0| + k^n |a_n|} \right) \right\} |P(z)|^2 \\ \leq (n \|P\| - |P'(z)|)^2. \end{aligned}$$

Equivalently for $|z| = 1$,

$$2n \|P\| |P'(z)| \leq n^2 \|P\|^2 - n \left\{ n \frac{k-1}{k+1} + \frac{2}{k+1} \frac{|a_0| - k^n |a_n|}{|a_0| + k^n |a_n|} \right\} |P(z)|^2,$$

which after simplification gives (2.1). For $k = 1$, the above inequality is trivially true for points on $|z| = 1$ for which $P(z) = 0$ by (1.2). This completes the proof of Theorem 2.1.

PROOF OF THEOREM 2.2. By hypothesis $P(z)$ has all its zeros in $|z| \geq k$, where $k \geq 1$. If $P(z)$ has a zero on $|z| = k$, then

$$m = \min_{|z|=k} |P(z)| = 0,$$

and Theorem 2.2 follows from Theorem 2.1 in this case. Henceforth, we suppose that all the zeros of $P(z)$ lie in $|z| > k$, $k \geq 1$. It follows by Rouché's theorem that for each β with $|\beta| < 1$, the polynomial $G(z) = P(z) - \beta m$, will have all its zeros in $|z| > k$. On applying inequality (4.2) to the polynomial $G(z) = P(z) - \beta m$, we get for every $|\beta| < 1$ and $|z| = 1$,

$$\begin{aligned} \left[|P'(z)|^2 + n \left\{ n \frac{k-1}{k+1} + \frac{2}{k+1} \left(\frac{|a_0 - \beta m| - k^n |a_n|}{|a_0 - \beta m| + k^n |a_n|} \right) \right\} |P(z) - \beta m|^2 \right]^{1/2} \\ \leq |Q'(z) - \bar{\beta} m n z^{n-1}|. \end{aligned} \quad (4.3)$$

By triangle inequality, we have

$$|P(z) - \beta m| \geq \left| |P(z)| - |\beta m| \right|, \quad (4.4)$$

and note that

$$\left| |P(z)| - |\beta m| \right|^2 = (|P(z)| - |\beta m|)^2, \quad (4.5)$$

so that, from (4.4) and (4.5), we obtain that

$$|P(z) - \beta m|^2 \geq (|P(z)| - |\beta|m)^2. \quad (4.6)$$

Also, all the zeros of $Q(z) = z^n \overline{P(1/\bar{z})}$ lie in $|z| \leq 1/k$, where $1/k \leq 1$, it follows by applying Lemma 3.4 to $Q(z)$, that for $|z| = 1$,

$$|Q'(z)| \geq nk^n \min_{|z|=1/k} |Q(z)|.$$

Since

$$\min_{|z|=1/k} |Q(z)| = \frac{1}{k^n} \min_{|z|=k} |P(z)| = \frac{m}{k^n},$$

therefore, we have

$$|Q'(z)| \geq mn. \quad (4.7)$$

From (4.7), it is possible to choose the argument of β on the right hand side of (4.3) such that for $|z| = 1$,

$$|Q'(z) - \overline{\beta} mn z^n| = |Q'(z)| - |\beta| mn. \quad (4.8)$$

Also, for every β , we have

$$|a_0 - \beta m| \geq |a_0| - |\beta|m,$$

and since the function

$$x \mapsto \frac{x - k^n |a_n|}{x + k^n |a_n|} \quad (x \geq 0)$$

is non-decreasing, it follows that

$$\frac{|a_0 - \beta m| - k^n |a_n|}{|a_0 - \beta m| + k^n |a_n|} \geq \frac{|a_0| - |\beta|m - k^n |a_n|}{|a_0| - |\beta|m + k^n |a_n|} = C_n(k, |\beta|), \quad (4.9)$$

where

$$C_n(k, t) = \frac{|a_0| - tm - k^n |a_n|}{|a_0| - tm + k^n |a_n|}.$$

Using (4.6), (4.8) and (4.9) in (4.3), we get for every $|\beta| < 1$ and $|z| = 1$,

$$\begin{aligned} \left\{ |P'(z)|^2 + n \left[n \frac{k-1}{k+1} + \frac{2}{k+1} C_n(k, |\beta|) \right] (|P(z)| - |\beta|m)^2 \right\}^{1/2} \\ \leq |Q'(z)| - |\beta| mn, \end{aligned}$$

which gives by using Lemma 3.5 for $|z| = 1$, that

$$|P'(z)|^2 + n \left[n \frac{k-1}{k+1} + \frac{2}{k+1} C_n(k, |\beta|) \right] (|P(z)| - |\beta|m)^2 \leq [n(\|P\| - |\beta|m) - |P'(z)|]^2.$$

Equivalently for $|z| = 1$,

$$2n(\|P\| - |\beta|m)|P'(z)| \leq n^2(\|P\| - |\beta|m)^2 - n \left[n \frac{k-1}{k+1} + \frac{2}{k+1} C_n(k, |\beta|) \right] (|P(z)| - |\beta|m)^2,$$

which gives for $|z| = 1$ and $|\beta| < 1$,

$$|P'(z)| \leq \frac{|P(z)| - |\beta|m}{2} \left\{ n - \left[\frac{(k-1)n}{k+1} + \frac{2C_n(k, |\beta|)}{k+1} \right] \left(\frac{|P(z)| - |\beta|m}{\|P\| - |\beta|m} \right)^2 \right\}.$$

Finally, for β with $|\beta| = 1$, the above inequality follows by continuity. Taking $|\beta| = t$, so that $0 \leq t \leq 1$, we get (2.4), and Theorem 2.2 is proved.

Acknowledgement. The work of the first author was supported in part by the Serbian Academy of Sciences and Arts (Φ -96).

REFERENCES

- [1] U. M. Ahangar, W. M. Shah, *Inequalities for the derivative of a polynomial with restricted zeros*, J. Anal. (2021), preprint.
- [2] A. Aziz, Q. M. Dawood, *Inequalities for a polynomial and its derivative*, J. Approx. Theory **54** (1988), 306–313.
- [3] A. Aziz, N. A. Rather, *New L^q inequalities for polynomials*, Math. Inequal. Appl. **1** (1998), 177–191.
- [4] S. Bernstein, *Sur l'ordre de la meilleure approximation des fonctions continues par des polynômes de degré donné*, Mem. Acad. R. Belg. **4** (1912), 1–103.
- [5] N. K. Govil, *Some inequalities for derivative of polynomials*, J. Approx. Theory **66** (1991), 29–35.
- [6] N. K. Govil, P. Kumar, *On sharpening of an inequality of Turán*, Appl. Anal. Discrete Math. **13** (2019), 711–720.

- [7] N. K. Govil, Q. I. Rahman, *Functions of exponential type not vanishing in a half plane and related polynomials*, Trans. Amer. Math. Soc. **137** (1969), 501–517.
- [8] P. Kumar, *On the inequalities concerning polynomials*, Complex Anal. Oper. Theory, **14** (2020), Art.65, pp. 1–11.
- [9] P. Kumar, R. Dhankhar, *Some refinements of inequalities for polynomials*, Bull. Math. Soc. Sci. Math. Roumanie **63**(111) (2020), 359–367.
- [10] P. D. Lax, *Proof of a conjecture of P. Erdős on the derivative of a polynomial*, Bull. Amer. Math. Soc. **50** (1944), 509–513.
- [11] M. A. Malik, *On the derivative of a polynomial*, J. London Math. Soc. **1** (1969), 57–60.
- [12] G. V. Milovanović, A. Mir, *On the Erdős-Lax inequality concerning polynomials*, Math. Inequal. Appl. **23** (2020), 1499–1508.
- [13] G. V. Milovanović, D. S. Mitrinović, Th. M. Rassias, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific, Singapore, 1994.
- [14] A. Mir, I. Hussain, *On the Erdős-Lax inequality concerning polynomials*, C. R. Acad. Sci. Paris Ser. I **355** (2017), 1055–1062.
- [15] A. Mir, D. Breaz, *Bernstein and Turán-type inequalities for a polynomial with constraints on its zeros*, RACSAM, **115** (2021), Art.124, pp. 1–12.
- [16] Q. I. Rahman, G. Schmeisser, *Analytic Theory of Polynomials*, Oxford University Press, 2002.

Serbian Academy of Sciences and Arts

11000 Beograd, Serbia

&

University of Niš

Faculty of Sciences and Mathematics

18000 Niš, Serbia

e-mail: `gvm@mi.sanu.ac.rs` (G. V. Milovanović)

Department of Mathematics

University of Kashmir

Srinagar, 190006, India

e-mails: `drabmir@yahoo.com` (A. Mir)

`malikadil6909@gmail.com` (A. Hussain)

`abrarahmad1100@gmail.com` (A. Ahmad)