



ACADEMIE SERBE DES SCIENCES ET DES ARTS

# BULLETIN

TOME CLI

CLASSE DES SCIENCES  
MATHEMATIQUES ET NATURELLES

SCIENCES MATHEMATIQUES

N° 43

# BULLETIN

TOME CLI

CLASSE DES SCIENCES  
MATHÉMATIQUES ET NATURELLES

SCIENCES MATHÉMATIQUES

N° 43





ISSN 0561-7332

ACADEMIE SERBE DES SCIENCES ET DES ARTS

---

---

# BULLETIN

TOME CLI

CLASSE DES SCIENCES  
MATHEMATIQUES ET NATURELLES

SCIENCES MATHEMATIQUES

N° 43

Rédacteur

GRADIMIR V. MILOVANOVIĆ

Membre de l'Académie

BEOGRAD 2018

Publie et imprimé par  
*Académie serbe des sciences et des arts*  
Beograd, Knez Mihailova 35

Tirage 300 exemplaires

© Académie serbe des sciences et des arts, 2018

## TABLE DES MATIÈRES

1. In Memorium: <i>Bogoljub Stanković (November 1, 1924 – May 16, 2018)</i> ...	1
2. I. Gutman, T. Réti: <i>Note on irregular graphs</i> .....	5
3. D. Cvetković, Z. Dražić, V. Kovačević-Vujčić, M. Čangalović: <i>The traveling salesman problem: The spectral radius and the length of an optimal tour</i> .....	17
4. T. Atanacković, S. Pilipović: <i>Bogoljub Stanković: Contributions to generalized asymptotics and mechanics</i> .....	27
5. G. V. Milovanović: <i>An efficient computation of parameters in the Rys quadrature formula</i> .....	39
6. M. Kostić: <i>Composition principles for generalized almost periodic functions</i> .....	65
7. N. Teofanov: <i>Wave-front sets in non-quasianalytic setting for Fourier- Lebesgue and modulation spaces</i> .....	81
8. Lj. Gajić, N. M. Ralević: <i>A common fixed point result in strong JS-metric space</i> .....	113

## A COMMON FIXED POINT RESULT IN STRONG JS-METRIC SPACE

LJILJANA GAJIĆ, NEBOJŠA RALEVIĆ

*Dedicated to Professor Bogoljub Stanković (1924–2018)*

(Presented at the 8th Meeting, held on November 28, 2018)

*A b s t r a c t.* The aim of our paper is the prove that the common fixed point result due to Sehgal and Thomas is valied in a class of generalized metric spaces in sence of Jleli and Samet.

AMS Mathematics Subject Classification (2000): 47H10, 54H25.

Key Words: common fixed point, JS-metric space, family of pointwise contractive self mappings.

### 1. Introduction

Metric fixed point theory, motivated by Banach fixed point theorem, is one of the most powerful and useful tools in nonlinear functional analysis. The application of this theory is remarkable in wide scale of mathematical, engineering, economic, physical, computer science and other fields of science. Following the Banach contraction principle in many papers the existence of weaker contractive conditions combined with stronger additional assumptions on the mapping or on the space, are investigated.

In [8] Sehgal initiated the study of fixed point for mappings with contractive iterate at a point. This result was extended by many authors (see [2], [3], [4], [5]).

**Definition 1.1.** [8] In metric space  $(X, d)$ , the mapping  $T : X \rightarrow X$  is said to be with contractive iterate at a point  $x \in X$  if there is a positive integer  $n(x)$  such that for all  $y \in X$

$$d(T^{n(x)}x, T^{n(x)}y) \leq qd(x, y), \quad (1.1)$$

where  $q \in (0, 1)$ .

In [9] V. M. Sehgal and J. W. Thomas proved a common fixed point result for a family of pointwise contractive self-mappings in metric space  $(X, d)$ .

**Definition 1.2.** [9] Let  $(X, d)$  be a complete metric space and  $M \subseteq X$ . Let  $\mathcal{F}$  be a commutative semigroup of self mappings (not necessarily continuous) of  $M$ . The semigroup  $\mathcal{F}$  is pointwise contractive in  $M$  if for each  $x \in M$ , there is an  $f_x \in \mathcal{F}$  such that

$$d(f_x(y), f_x(x)) \leq \varphi(d(y, x)), \quad (1.2)$$

for all  $y \in M$ , where  $\varphi$  is some real valued function defined on the nonnegative reals.

**Theorem 1.1.** [9] Let  $M$  be a closed subset of  $X$  and  $\mathcal{F}$  a commutative semigroup of self-mappings of  $M$ , which is pointwise contractive in  $M$  for some  $\varphi : [0, \infty) \rightarrow [0, \infty)$ , where  $\varphi$  is nondecreasing, continuous on the right and satisfies  $\varphi(r) < r$  for all  $r > 0$ . If for some  $x_0 \in M$ ,

$$\sup\{d(f(x_0), x_0) : f \in \mathcal{F}\} < \infty, \quad (1.3)$$

then, there exists a unique  $\xi \in M$  such that  $f(\xi) = \xi$  for each  $f \in \mathcal{F}$ . Moreover, there is a sequence  $\{g_n\} \subset \mathcal{F}$  with  $g_n(x) \rightarrow \xi$  for each  $x \in M$ .

On the other side, recent research in fixed point theory are focused on generalizing the underlying space. Consideration various generalization of metric spaces (partial metric spaces, fuzzy metric spaces, probability metric spaces, quasi-metric spaces, uniform spaces, ultra metric spaces,  $b$ -metric spaces, cone metric spaces and so on) leads to opportunity to use distinct advantages by creating topological structure suitable for application in some cases when the classical metric does not give the answer.

Recently, Jleli and Samet [6] introduced a generalization of the notion of a metric spaces which they called a generalized metric space. They noted that some above-mentioned abstract metric spaces may be regarded as particular cases of their general definition. They also stated and proved fixed point theorems for some contractions defined on these spaces.

In this paper we consider Sehgal and Thomas common fixed point result in setting of generalized metric spaces in sense of Jleli and Samet.



### 2. Preliminaries

Let  $X$  be a nonempty set and  $\mathcal{D} : X \times X \rightarrow [0, \infty]$  be a mapping. For every  $x \in X$  define a set

$$C(\mathcal{D}, X, x) = \{\{x_n\} \subset X : \lim_{n \rightarrow \infty} \mathcal{D}(x_n, x) = 0\}.$$

In this case, we say that  $\mathcal{D}$  is a generalized metric in the sense of Jleli and Samet [6] (for short JS-metric) on  $X$  if it satisfies the following conditions:

- ( $\mathcal{D}_1$ ) for every  $(x, y) \in X \times X$ ,  $\mathcal{D}(x, y) = 0 \Rightarrow x = y$ ;
- ( $\mathcal{D}_2$ ) for every  $(x, y) \in X \times X$ ,  $\mathcal{D}(x, y) = \mathcal{D}(y, x)$ ;
- ( $\mathcal{D}_3$ ) there exists  $C > 0$  such that for every  $(x, y) \in X \times X$  and  $\{x_n\} \in C(\mathcal{D}, X, x)$ ,

$$\mathcal{D}(x, y) \leq C \limsup_{n \rightarrow \infty} \mathcal{D}(x_n, y).$$

In this case  $(X, \mathcal{D})$  is said to be JS-metric space. Note that, if  $C(\mathcal{D}, X, x) = \emptyset$  for all  $x \in X$ , then ( $\mathcal{D}_3$ ) is trivially hold.

The class of JS-metric spaces is larger than many known class of abstract metric spaces. For example, every standard metric space, every  $b$ -metric space, every dislocated metric space (in the sense of Hitzler-Seda), and every modular space with the Fatou property is a JS-metric space. For more details see [6].

Given a JS-metric space  $(X, \mathcal{D})$  and a point  $x \in X$ , a sequence  $\{x_n\} \subseteq X$  is said to be:

1°  $\mathcal{D}$  – convergent to  $x$  if  $\{x_n\} \in C(\mathcal{D}, X, x)$ , in such a case, we will write

$$\{x_n\} \xrightarrow{\mathcal{D}} x;$$

2°  $\mathcal{D}$  – Cauchy if

$$\lim_{n, m \rightarrow \infty} \mathcal{D}(x_n, x_m) = 0. \tag{2.1}$$

A JS-metric space  $(X, \mathcal{D})$  is *complete* if every  $\mathcal{D}$ -Cauchy sequence in  $X$  is  $\mathcal{D}$ -convergent.

Jleli and Samet proved that the limit of a  $\mathcal{D}$ -convergent sequence is unique.

**Proposition 2.1.** [6] *Let  $(X, \mathcal{D})$  be a JS-metric space. Let  $\{x_n\}$  be a sequence in  $X$  and  $(x, y) \in X \times X$ . If  $\{x_n\}$   $\mathcal{D}$ -converges to  $x$  and  $\{x_n\}$   $\mathcal{D}$ -converges to  $y$ , then  $x = y$ .*

They also proved fixed point theorem for  $q$ -contraction defined on these spaces as a generalization of Banach contraction principle.

**Theorem 2.1.** [6] *Suppose that the following conditions hold:*

- (i)  $(X, \mathcal{D})$  is a complete JS-metric space,
- (ii)  $f : X \rightarrow X$  is a  $k$ -contraction for some  $k \in (0, 1)$ , that is,

$$\mathcal{D}(f(x), f(y)) \leq k\mathcal{D}(x, y)$$

for all  $(x, y) \in X \times X$ .

- (iii) there exists  $x_0 \in X$  such that  $\sup\{\mathcal{D}(x_0, f^n(x_0)) : n \in \mathbb{N}\} < \infty$ .

Then  $\{f^n(x_0)\}$  converges to  $\omega \in X$ , a fixed point of  $f$ . Moreover, if  $\omega' \in X$  is another fixed point of  $f$  such that  $\mathcal{D}(\omega, \omega') < \infty$ , then  $\omega = \omega'$ .

For more fixed point results in JS-metric spaces we refer the reader to [6], [7].

Now we are going to introduce an interesting subclass of JS-metric spaces as a frame for our researches.

**Definition 2.1.** Let  $X$  be a nonempty set and let  $\mathcal{D}^* : X \times X \rightarrow [0, \infty)$  be a function which satisfies the following conditions:

- ( $\mathcal{D}^*1$ )  $\mathcal{D}^*(x, y) = 0 \Leftrightarrow x = y$ ;
- ( $\mathcal{D}^*2$ )  $\mathcal{D}^*(x, y) = \mathcal{D}^*(y, x)$ , for all  $x, y \in X$ ;
- ( $\mathcal{D}^*3$ ) there exists  $C > 0$  such that,  
for every  $x, y \in X$  and  $\{x_n\} \in C(\mathcal{D}^*, X, x)$ ,  $\{y_n\} \in C(\mathcal{D}^*, X, y)$   
 $\mathcal{D}^*(x, y) \leq C \limsup_{n \rightarrow \infty} \mathcal{D}^*(x_n, y_n)$ .

Then  $\mathcal{D}^*$  is a strong JS-metric and the pair  $(X, \mathcal{D}^*)$  is a strong JS-metric space.

**Remark 2.1.** Since sequence  $x_n = x$ ,  $n \in \mathbb{N}$ ,  $\mathcal{D}^*$ -converges to  $x$  every strong JS-metric space is JS-metric space.

**Example 2.1.** Every standard metric space  $(X, d)$  is a strong JS-metric space.

Since, we only have to check condition ( $\mathcal{D}^*3$ ), let  $x, y \in X$ ,  $\{x_n\} \xrightarrow{d} x$ , and  $\{y_n\} \xrightarrow{d} y$ . Using triangle inequality twice

$$d(x, y) \leq d(x, x_n) + d(x_n, y_n) + d(y_n, y)$$

for all naturale number  $n$ , so

$$d(x, y) \leq \limsup_{n \rightarrow \infty} d(x_n, y_n)$$

and ( $\mathcal{D}^*3$ ) is satisfied with  $C = 1$ .

**Example 2.2.** Every  $b$ -metric space  $(X, d_b)$  is a strong metric space.

Let us recall the definition of  $b$ -metric space.

Let  $d_b : X \times X \rightarrow (0, +\infty)$  be a given mapping. We say that  $d_b$  is a  $b$ -metric on  $X$  if it satisfies the following conditions:

(b1) for every  $(x, y) \in X \times X$ , we have

$$d_b(x, y) = 0 \Leftrightarrow x = y;$$

(b2) for every  $(x, y) \in X \times X$ ,

$$d_b(x, y) = d_b(y, x);$$

(b3) there exists  $s \geq 1$  such that, for every  $x, y, z \in X$ ,

$$d_b(x, y) \leq s(d_b(x, z) + d_b(z, y)).$$

In this case,  $(X, d_b)$  is said to be a  $b$ -metric space.

Obviously, we have to prove that  $d_b$  satisfies the property  $(D^*3)$ . Let  $x, y \in X$  and  $\{x_n\} \xrightarrow{d_b} x, \{y_n\} \xrightarrow{d_b} y$ . By property (b3) we have that

$$\begin{aligned} d_b(x, y) &\leq s(d_b(x, x_n) + d_b(x_n, y)) \\ &\leq s d_b(x, x_n) + s^2 d_b(x_n, y_n) + s^2 d_b(y_n, y) \end{aligned}$$

for every natural number  $n$ . Thus, we have

$$d_b(x, y) \leq s^2 \limsup_{n \rightarrow \infty} d_b(x_n, y_n)$$

and the property  $(D^*3)$  is satisfied with  $c = s^2$ .

### 3. Main result

A generalization of the contraction principle can be obtained using different type of a nondecreasing function  $\varphi : [0, \infty) \rightarrow [0, \infty)$ . The most usual additional properties imposed on  $\varphi$  are:

$$(\varphi_1) \varphi(0) = 0,$$

$$(\varphi_2) \varphi \text{ is right continuous and } \varphi(t) < t, \text{ for all } t > 0,$$

$$(\varphi_3) \lim_{i \rightarrow \infty} \varphi^i(t) = 0, \text{ for all } t > 0,$$

$$(\varphi_4) \{t_i\} \subset [0, \infty) \text{ is a sequence such that } t_{i+1} \leq \varphi(t_i), \text{ then } \lim_{i \rightarrow \infty} t_i = 0.$$

It is well known that

$$(\varphi_4) \Leftrightarrow (\varphi_3) \Leftrightarrow (\varphi_2) \Rightarrow (\varphi_1),$$

and  $\varphi$  with property  $(\varphi_2)$ , is continuous at 0.

**Theorem 3.1.** *Let  $B$  be a closed nonempty subset of complete strong JS-metric space  $(X, \mathcal{D}^*)$  and  $\mathcal{F}$  a commutative semigroup of self-mapping of  $B$  pointwise contractive in  $B$  for some  $\varphi : [0, \infty) \rightarrow [0, \infty)$ , where  $\varphi$  is nondecreasing continuous on the right and satisfies  $\varphi(t) < t$ , for all  $t > 0$ .*

*If for some  $x_0 \in B$*

$$\sup\{\mathcal{D}^*(x_0, f(x_0)) : f \in \mathcal{F}\} < \infty$$

*then, there exists a unique  $u \in B$  such that  $f(u) = u$  for each  $f \in \mathcal{F}$ . Moreover, there is a sequence  $\{g_n\} \subset \mathcal{F}$  with  $\{g_n(x)\} \xrightarrow{\mathcal{D}^*} u$ , for each  $x \in B$ .*

PROOF. If  $d_0 = \sup\{\mathcal{D}^*(x_0, f(x_0)) : f \in \mathcal{F}\}$ , then  $\varphi^n(d_0) \rightarrow 0$ ,  $n \rightarrow \infty$ . Let  $f_0 = f_{x_0}$  and inductively  $f_n = f_{x_n}$ , where  $x_{n+1} = f_n(x_n)$ . Then, for a fixed  $k \geq 0$

$$\sup_{n \geq k} \mathcal{D}^*(x_{n+1}, x_{k+1}) = \sup_{n \geq k} \mathcal{D}^*(f_n \circ f_{n-1} \circ \cdots \circ f_k(x_k), f_k(x_k)).$$

Let  $h_n = f_n \circ f_{n-1} \circ \cdots \circ f_{k+1}$ . It follows that

$$\begin{aligned} \sup_{n \geq k} \mathcal{D}^*(x_{n+1}, x_{k+1}) &= \sup_{n \geq k} \mathcal{D}^*(f_k(h_n(x_k)), f_k(x_k)) \\ &\leq \sup_{n \geq k} \varphi(\mathcal{D}^*(h_n(x_k), x_k)) \\ &\leq \sup_{n \geq k} \varphi^{k+1}(\mathcal{D}^*(h_n(x_0), x_0)) \rightarrow 0, \end{aligned}$$

when  $k \rightarrow \infty$ . Thus, the sequence  $\{x_n\}$  is  $\mathcal{D}^*$ -Cauchy. Let  $\{x_n\} \xrightarrow{\mathcal{D}^*} u \in B$ . By hypotheses, there is a  $f_u \in \mathcal{F}$  such that

$$\mathcal{D}^*(f_u(x_n), f_u(u)) \leq \varphi(\mathcal{D}^*(x_n, u)) \rightarrow 0, \quad n \rightarrow \infty,$$

so that

$$\{f_u(x_n)\} \xrightarrow{\mathcal{D}^*} f_u(u).$$

On the other side  $\mathcal{D}^*(x_k, f_u(x_k)) \leq \varphi^k(\mathcal{D}^*(x_0, f_u(x_0))) \rightarrow 0$ ,  $k \rightarrow \infty$ , so that

$$\mathcal{D}^*(u, f_u(u)) \leq C \cdot \limsup_{n \rightarrow \infty} \mathcal{D}^*(x_k, f_u(x_k)) \rightarrow 0, \quad k \rightarrow \infty,$$

and consequently  $f_u(u) = u$ .

Using (1.2) one can prove that  $u$  is a unique fixed point of  $f_u$  on  $B$ . Furthermore, since  $\mathcal{F}$  is commutative, for any  $f \in \mathcal{F}$

$$f(u) = f(f_u(u)) = f_u(f(u))$$

and therefore  $f(u) = u$ , for all  $f \in \mathcal{F}$ .

For each nonnegative integer  $n$ , set  $g_n = f_u^n$ . Obviously  $g_n \in \mathcal{F}$ .

For any fixed  $x \in B$

$$\mathcal{D}^*(g_n(x), u) = \mathcal{D}^*(f_u^n(x), f_u(u)) \leq \varphi^n(\mathcal{D}^*(x, u)) \rightarrow 0, \quad n \rightarrow \infty,$$

so  $g_n(x) \xrightarrow{\mathcal{D}^*} u$ , and proof is completed.

**Corollary 3.1.** *Let  $B$  be a closed subset of complete strong JS-metric space  $(X, \mathcal{D}^*)$  and  $f$  a self-mapping of  $B$ . If  $f$  satisfies conditions:*

1° *there exists  $x_0 \in B$  such that  $\sup\{\mathcal{D}^*(x_0, f^k(x_0)) : k \in \mathbb{N}\} < \infty$ ;*

2° *for each  $x \in B$ , there exists an integer  $n(x) \geq 1$  such that for all  $y \in B$*

$$\mathcal{D}^*(f^{n(x)}(x), f^{n(x)}(y)) \leq \varphi(\mathcal{D}^*(x, y)),$$

where  $\varphi : [0, \infty) \rightarrow [0, \infty)$  is nondecreasing right continuous function and  $\varphi(t) < t$ , for  $t > 0$ , then there exists a unique  $u \in B$  such that  $f(u) = u$  and  $\{f^k(x)\} \xrightarrow{\mathcal{D}^*} u$ , for any  $x \in B$ .

PROOF. Family  $\mathcal{F} = \{f^k : k \in \mathbb{N}\}$  is a commutative semigroup of pointwise contractive mappings in  $B$  so by Theorem 3.1, there exists a unique fixed point  $u$  of  $f$  and there is  $\{g_k\} \subset \mathcal{F}$  such that  $\{g_k(x)\} \xrightarrow{\mathcal{D}^*} u$ , for each  $x \in B$ .

Let us prove that  $\{f^k(x)\} \xrightarrow{\mathcal{D}^*} u$ , for any  $x \in B$ . For  $k$  sufficiently large we have  $k = r \cdot n(u) + s$ , with  $r > 0$  and  $0 \leq s < n(u)$  and therefore for any  $x \in B$

$$\begin{aligned} \mathcal{D}^*(f^k(x), u) &= \mathcal{D}^*(f^{r \cdot n(u) + s}(x), f^{n(u)}(u)) \\ &\leq \varphi^r(\mathcal{D}^*(f^s(x), u)) \\ &\leq \varphi^r(d_0) \end{aligned}$$

for  $d_0 = \max\{\mathcal{D}^*(f^s(x), u) : 0 \leq s < n(u)\}$ . Since  $\varphi^r(d_0) \rightarrow 0$ ,  $r \rightarrow \infty$ , it follows that  $\{f^k(x)\} \xrightarrow{\mathcal{D}^*} u$ , for all  $x \in B$ , so proof is completed.

The next corollary is a generalization of Guseman fixed point result from [5].

**Corollary 3.2.** *Let  $B$  be a closed subset of complete strong JS-metric space  $(X, \mathcal{D}^*)$  and  $f$  a self-mapping of  $B$ . If  $f$  is a mapping with contractive iterate at any point  $x \in B$  for some  $0 < q < 1$  and there exists  $x_0 \in B$  such that  $\sup\{\mathcal{D}^*(x_0, f^k(x_0)) : k \in \mathbb{N}\} < \infty$ , then there exists a unique  $u \in B$  such that  $f(u) = u$  and  $\{f^k(x)\} \xrightarrow{\mathcal{D}^*} u$  for any  $x \in B$ .*

PROOF. Take,  $\varphi(t) = qt, t \in [0, \infty)$ , and use Corollary 3.1.

**Acknowledgement.** This research is supported by grants 174024 and 174009 of Ministry of Education, Science and Technological Development, Republic of Serbia.

#### REFERENCES

- [1] Y. ElKouch, El M. Marhrani, *On some fixed point theorems in generalized metric spaces*, Fixed Point Theory Appl. 2017, Paper No. 23, 17 pp.
- [2] Lj. Gajić, Z. Lozanov-Crvenković, *On mappings with contractive iterate at a point in generalized metric spaces*, Fixed Point Theory Appl. 2010, Art. ID 458086, 16 pp.
- [3] Lj. Gajić, M. Stojaković, *On Ćirić generalization of mappings with a contractive iterate at a point in  $G$ -metric space*, Appl. Math. Comput., **219** (2012), 435–441.
- [4] Lj. Gajić, M. Stojaković, *Sehgal-Thomas type fixed point theorems in generalized metric spaces*, Filomat, **31**:11 (2017), 3335–3346.
- [5] L. F. Guseman, *Fixed point theorems for mappings with a contractive iterate at a point*, Proc. Amer. Math. Soc., **26** (1970), 615–618.
- [6] M. Jleli, B. Samet, *A generalilzed metric space and related fixed point theorems*, Fixed Point Theory and Applications 2015, (2015), 14 pages.
- [7] E. Karapinar, D. O'Regan, A. F. Roldán López de Hierro, N. Shahzad, *Fixed point theorems in new generalized metric spaces*. J. Fixed Point Theory Appl. **18** (2016), no. 3, 645–671.
- [8] V. M. Sehgal, *A fixed point theorem for mappings with a contractive iterate*, Proc. Amer. Math. Soc., **23** (1969), 631–634.
- [9] V. M. Sehgal, J.W. Thomas, *A fixed point theorem for semigroups of mappings with a contractive iterate*, Proc. Amer. Math. Soc., **29** (3) (1971), 565–570.

Faculty of Science  
University of Novi Sad  
Trg Dositeja Obradovića 4  
Novi Sad 21125, Serbia  
e-mail: [gajic@dmi.uns.ac.rs](mailto:gajic@dmi.uns.ac.rs)

Faculty of Technical Sciences  
University of Novi Sad  
Trg Dositeja Obradovića 6  
Novi Sad 21125, Serbia  
e-mail: [nralevic@uns.ac.rs](mailto:nralevic@uns.ac.rs)